













# My vaccinations

The Child and Family Agency recommends following **the full basic schedule** suggested by the Superior Health Council. Most vaccines are **free**. The **polio vaccination is required by law**. Your child can be vaccinated **at a Child and Family Agency health clinic**, at the family doctor's office or at the paediatrician's office. The doctor or nurse will note here and in the online record which vaccines your child has received and when.

Vaccination against	8 weeks	12 weeks	16 weeks	12 months	15 months
 <b>COMBINATION VACCINE</b> <b>polio</b> (infantile paralysis) <b>diphtheria</b> (croup) <b>tetanus</b> (lockjaw) <b>pertussis</b> (whooping cough) <b>hepatitis B</b> (jaundice) <b>haemophilus influenzae type b</b> (meningitis)					
 <b>pneumococci</b> (meningitis + pneumonia) At a gestational age of <b>fewer than 37 weeks</b> , an <b>additional dose</b> will be administered <b>at 12 weeks of age</b> .		Extra dose for premature infants			
 <b>rotavirus</b> (gastroenteritis) The rotavirus vaccine is administered orally and costs about <b>€12 on a doctor's prescription</b> . Store in the refrigerator (2-8°C). It takes 2 or 3 doses, depending on the brand.					
 <b>COMBINATION VACCINE</b> <b>polio</b> (infantile paralysis) <b>diphtheria</b> (croup) <b>tetanus</b> (lockjaw) <b>pertussis</b> (whooping cough) <b>hepatitis B</b> (jaundice) <b>haemophilus influenzae type b</b> (meningitis)					
 <b>pneumococci</b> (meningitis + pneumonia)					
 <b>COMBINATION VACCINE</b> <b>measles</b> <b>mumps</b> (thick ear) <b>rubella</b> (German measles)					
 <b>meningococcal ACWY</b> (meningitis)					

 Only vaccines with this icon need to be paid for. All the others are free.

 Not applicable at this age.