My vaccinations

The Child and Family Agency recommends following **the full basic schedule** suggested by the Superior Health Council. Most vaccines are **free**. The **polio vaccination** is **required by law**. Your child can be vaccinated **at a Child and Family Agency health clinic**, at the family doctor's office or at the paediatrician's office. The doctor or nurse will note here and in the online record which vaccines your child has received and when.

Vaccination against	8 weeks	12 weeks	16 weeks		Vaccination against	12 months	IS INCIALIS If the gestational age is fewer than 37 weeks , vaccines are advanced from 15 months to 13 months.
COMBINATION VACCINE polio (infantile paralysis) diphtheria (croup) tetanus (lockjaw) pertussis (whooping cough) hepatitis B (jaundice) haemophilus influenzae type b (meningitis)				<u></u>	COMBINATION VACCINE polio (infantile paralysis) diphtheria (croup) tetanus (lockjaw) pertussis (whooping cough) hepatitis B (jaundice) haemophilus influenzae type b (meningitis)		
Pneumococci (meningitis + pneumonia)At a gestational age of fewer than 37 weeks, an additional dose will be administered at 12 weeks of age.		Extra dose for premature infants		jî) jî)	pneumococci (meningitis + pneumonia) COMBINATION VACCINE measles mumps (thick ear) rubella (German measles)		
rotavirus (gastroenteritis) The rotavirus vaccine is administered orally and costs about €12 on a doctor's prescription. Store in the refrigerator (2-8°C). It takes 2 or 3 doses, depending on the brand.	E	E	Extra dose when using Rota Teq ®	ſĮ.	meningococcal ACWY (meningitis)		
Conly vac	cines with this ic	on need to be paid	d for. All the others a	re free.	Not applicable at this age.		



15 months

彩